# PRINTER RUSH (PTO ASSISTANCE)

	Application :	09/97/11)	Examiner:	Pryor	GAU:	<u>1616</u> 04·23-05	
	From:		Location: (	FMF FDC	Date:	04.23-05	
Tracking #: <u>0(0)737(3</u> Week Date: <u>0(3)705</u>							
		DOC CODE  1449  1DS CLM IIFW SRFW DRW OATH 312 SPEC	2/162	MISCELL Continuing Foreign Price Document I Fees Other	Data ority		
	RUSH] MESSAGE: Aciples G7 +68 Contain a Chemical frombanitarial legisle data. Please Resolu. Thank Yw. P						
			- :				
[XRUSH] RESPONSE:  OVVEC TO  OVVEC T							
	NOTE: Thi	NOTE: This form will be included as part of the official USPTO record, with the Response document coded as XRUSH.					
	REV 10/04	oded as XRUSH. By monolly		5	P/ 360	-22	

# RECEIVED

## MAY 02 RECT

rublishing Division 13

P/3610-22

#### IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of

Anne-Marie CAMINADE et al.

Serial No.: 09/936,119

Filed: February 1, 2002

Date: May 2, 2005

Group Art Unit: 1616

Examiner: Pryor, A.N.

For: PESTICIDE AND/OR GROWTH REGULATING COMPOSITIONS

Commissioner for Patents P.O. Box 1450 Alexandria, Virginia 22313-1450

### TRANSMITTAL LETTER

Sir:

In accordance with the telephone conversation with Examiner Small on Friday, April 29, 2005, enclosed are pages 67 and 68 of the above-identified specification.

No new matter is added.

Respectfully submitted,

5 MAY 2005

Date

Paul Grandinetti


Reg. No. 30,754

OSTROLENK, FABER, GERB & SOFFEN, LLP

1180 Avenue of the Americas

New York, New York 10036-8403

Telephone (202) 429-4560



To facilitate understanding of the remainder of the present disclosure, the dendrimer represented by Figure (X) and whose terminal functional groups comprise chemical radicals derived from Girard T reagents is called dendrimer G'4-T.

For the preparation of said dendrimer G'4-T, a dendrimer termed G'4-CHO is normally used whose

aldehyde type groups at the periphery, preferably all said terminal functional groups consist of aldehyde-type groups at the periphery; said dendrimer G'4-CHO may be prepared with reference to the information given in the manual <u>Les dendrimères</u> previously cited in the present disclosure. For the preparation of said dendrimer G'4-CHO, the reaction scheme represented by Figure (XI) below may be followed for example.

$$S = PCI_{3}$$

$$S = PCI_{3}$$

$$S = P - (O - CHO)_{3}$$

$$S = P - (O - CHO)_{4}$$

$$S = P - (O - CHO)_{5}$$

It is then possible to react said dendrimer G'4-CHO in the presence of the so-called Girard T reagent as described above in the present text and a representation of which is given below by Figure (V), and thus to obtain the dendrimer termed G'4-T.